

16% Phoenix HEIFER GROWER- R Medicated

Medicated Calf Feed. Medicated for the prevention and control of coccidiosis caused by Eimeria bovis and Eimeria zuernii.

ACTIVE DRUG INGREDIENT

Monensin.....40 g/ton

GUARANTEED ANALYSIS

Crude Protein (Minimum).....	16.00%
Crude Fat (Minimum).....	2.40%
Crude Fiber (Maximum).....	7.00%
ADF (Maximum).....	8.60%
Calcium (Minimum).....	0.66%
Calcium (Maximum).....	0.98%
Phosphorus (Minimum).....	0.40%
Selenium (Minimum).....	.30ppm
Vitamin A.....	11,000 IU/LB

INGREDIENTS

Grain Products, Processed Grain By-products, Plant Protein Products, Calcium Carbonate, Milk Protein Product, Roughage Products, Sodium Sesquicarbonate, Potassium Chloride, Sodium Chloride, Vitamin A Supplement, Carbohydrate (Glucose or corn starch), Gelatin, Ethoxyquine, Cane Molasses, Liquid Feed Syrup, Vegetable Fat (Feed Grade), Phosphoric acid, Soy Lecithin, Sulfuric Acid, Propionic Acid (A Preservative), Sorbic Acid (A Preservative), Benzoic Acid (A Preservative), Sodium Sulfate, Sodium Sulfite, Potassium Sulfate, Sodium Bentonite, Vitamin E Supplement, Manganese Sulfate, Zinc Sulfate, Copper Sulfate, Mineral Oil, Magnesium Oxide, Vitamin D3 Supplement, Cobalt Sulfate, Montmorillonite Clay, Diatomaceous Earth Ethylenediamine Dihydroiodide, Sodium Selenite, Yeast Brewers Dehydrated, Yeast Extract, Calcium Propionate (a preservative), Magnesium Silicate, BHA (a preservative-antioxidants), Natural and Artificial Flavoring.

FEEDING DIRECTIONS

Feed continuously at the rate of 1.25 pound of feed/100 pound of body weight; do not exceed 10.00 pounds per day. Feeding at 1.25 pound of feed/100 pound of body weight will supply Monensin at the level to control coccidiosis. Feed 16% Phoenix Heifer Grower - R daily and supply fresh water at all times. Top quality hay may be introduced once dry feed intake reaches 5 pounds per day.

Caution: Do not allow horses or other equines access to formulations containing Rumensin. Ingestion of Rumensin by equines has been fatal. Rumensin medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Rumensin has been fatal to cattle and could be fatal in goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Rumensin recommended in the feeding directions, as reduced average daily gains may result.

Warning: A withdrawal period has not been established for Monensin in pre-ruminating calves. Do not use in calves to be processed for veal.

03.07.24
Manufactured For
Phoenix Feeds, New Haven, VT 05472

Net Weight 50 pounds

16% Phoenix HEIFER GROWER- R w/Clarifly® Larvicide Medicated

Medicated Calf Feed. Medicated for the prevention and control of coccidiosis caused by Eimeria bovis and Eimeria zuernii. Diflubenzuron added as an insect growth regulator, which prevents the development of house, stable, face and horn flies in the manure of treated calves.

ACTIVE DRUG INGREDIENT

Monensin.....40 g/ton
Diflubenzuron..... 3.60 g/ton

GUARANTEED ANALYSIS

Crude Protein (Minimum)..... 16.00%
Crude Fat (Minimum)..... 2.60%
Crude Fiber (Maximum)..... 4.95%
ADF (Maximum)..... 7.30%
Calcium (Minimum)..... 0.70%
Calcium (Maximum)..... 0.79%
Phosphorus (Minimum)..... .39%
Selenium (Minimum)..... .30ppm
Vitamin A..... 13,000 IU/LB

INGREDIENTS

Grain Products, Processed Grain By-products, Plant Protein Products, Calcium Carbonate, Milk Protein Product, Roughage Products, Sodium Sesquicarbonate, Potassium Chloride, Sodium Chloride and Processed Grain By-Products, Calcium Carbonate and Vitamin A Supplement, Carbohydrate (Glucose or corn starch), Gelatin, Ethoxyquine, Cane Molasses, Liquid Feed Syrup, Vegetable Fat (Feed Grade), Phosphoric acid, Soy Lecithin, Sulfuric Acid, Propionic Acid (A Preservative), Sorbic Acid (A Preservative), Benzoic Acid (A Preservative), Monocalcium Phosphate, Dicalcium Phosphate, Vitamin E Supplement, Selenium Yeast, Calcium Sulfate, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper, Amino Acid Complex, Cobalt Glucoheptonate complex, Magnesium Oxide, Manganese Sulfate, Zinc Sulfate, Potassium Chloride, Vitamin A Supplement, Vitamin D3 Supplement, Copper Sulfate, Ethylenediamine Dihydrochloride, Cobalt Carbonate, Diflubenzuron (CAS#35367-38-5), Montmorillonite Clay, Diatomaceous Earth, Hydrated Sodium Calcium Aluminosilicate, Yeast Brewers Dehydrated, Yeast Extract, Calcium Propionate (a preservative), Citric Acid (a preservative), BHT (a preservative), Silicon Dioxide, Vegetable Oil, Ammonium Hydroxide, Water, Natural and Artificial Flavoring.

FEEDING DIRECTIONS

Feed continuously at the rate of 1.25 pound of feed/100 pound of body weight; do not exceed 10.00 pounds per day. Feeding at 1.25 pound of feed/100 pound of body weight will supply Monensin at the level to control coccidiosis. Feed 16% Phoenix Heifer Grower - R daily and supply fresh water at all times. Top quality hay may be introduced once dry feed intake reaches 5 pounds per day. When feed as directed this feed will provide sufficient Diflubenzuron Larvicide to prevent the emergence of adult house, stable, horn and face flies from the manure of treated cattle. Existing adult flies will not be affected. For optimal fly growth regulation, Diflubenzuron treated feed should first be fed 30 days prior to the last frost, continuously through the fly season and end 30 days after the first frost.

Caution: Do not allow horses or other equines access to formulations containing Rumensin. Ingestion of Rumensin by equines has been fatal. Rumensin medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Rumensin has been fatal to cattle and could be fatal in goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Rumensin recommended in the feeding directions, as reduced average daily gains may result.

Warning: A withdrawal period has not been established for Monensin in pre-ruminating calves. Do not use in calves to be processed for veal.

03.07.24
Manufactured For
Phoenix Feeds, New Haven, VT 05472

Net Weight 50 pounds