

18% Phoenix Calf Starter – R

Medicated

Medicated Calf Feed. Medicated for the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENT

Monensin.....40 g/ton

GUARANTEED ANALYSIS

Crude Protein (Minimum).....	18.00%
Crude Fat (Minimum).....	2.50%
Crude Fiber (Maximum).....	3.85%
ADF (Maximum).....	6.00%
Calcium (Minimum).....	0.80%
Calcium (Maximum).....	0.89%
Phosphorus (Minimum).....	0.42%
Selenium (Minimum).....	.30ppm
Vitamin A.....	12,000 IU/LB

INGREDIENTS

Grain Products, Processed Grain By-products, Plant Protein Products, Calcium Carbonate, Milk Protein Product, Roughage Products, Sodium Sesquicarbonate, Potassium Chloride, Sodium Chloride and Processed Grain By-Products, Calcium Carbonate and Vitamin A Supplement, Carbohydrate (Glucose or corn starch), Gelatin, Ethoxyquine, Cane Molasses, Liquid Feed Syrup, Vegetable Fat (Feed Grade), Phosphoric acid, Soy Lecithin, Sulfuric Acid, Propionic Acid (A Preservative), Sorbic Acid (A Preservative), Benzoic Acid (A Preservative), Monocalcium Phosphate, Dicalcium Phosphate, Vitamin E Supplement, Selenium Yeast, Calcium Sulfate, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper, Amino Acid Complex, Cobalt Glucoheptonate complex, Magnesium Oxide, Manganese Sulfate, Zinc Sulfate, Potassium Chloride, Vitamin A Supplement, Vitamin D3 Supplement, Copper Sulfate, Ethylenediamine Dihydroid, Cobalt Carbonate, Montmorillonite Clay, Diatomaceous Earth, Hydrated Sodium Calcium Aluminosilicate, Yeast Brewers Dehydrated, Yeast Extract, Calcium Propionate (a preservative), Citric Acid (a preservative), BHT (a preservative), Silicon Dioxide, Vegetable Oil, Ammonium Hydroxide, Water, Natural and Artificial Flavoring.

FEEDING DIRECTIONS

Feed continuously at the rate of 1.25 pound of feed/100 pound of body weight; do not exceed 10.0 pounds per day. Feeding at 1.25 pound of feed/100 pound of body weight will supply Monensin at the level to control coccidiosis. Feed 18% Phoenix Feeds Calf Starter - R daily and supply fresh water at all times. Top quality hay may be introduced once dry feed intake reaches 5 pounds per day.

Caution: Do not allow horses or other equine access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin-medicated cattle and goat feeds are safe for use in cattle goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the fed refusals and the amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning: A withdrawal period has not been established for Monensin in pre-ruminating calves. Do not use in calves to be processed for veal.

03.07.24

Manufactured For
Phoenix Feeds, New Haven, VT 05472

Net Weight 50 pounds

18% Phoenix Calf Starter – R w/Clarifly® Larvicide

Medicated

Medicated Calf Feed. Medicated for the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*. Diflubenzuron added as an insect growth regulator, which prevents the development of house, stable, face and horn flies in the manure of treated calves.

ACTIVE DRUG INGREDIENT

Monensin.....	40 g/ton
Diflubenzuron.....	3.60 g/ton

GUARANTEED ANALYSIS

Crude Protein (Minimum).....	18.00%
Crude Fat (Minimum).....	2.50%
Crude Fiber (Maximum).....	3.85%
ADF (Maximum).....	6.00%
Calcium (Minimum).....	0.80%
Calcium (Maximum).....	0.89%
Phosphorus (Minimum).....	0.42%
Selenium (Minimum).....	0.30ppm
Vitamin A.....	12,000 IU/LB

INGREDIENTS

Grain Products, Processed Grain By-products, Plant Protein Products, Calcium Carbonate, Milk Protein Product, Roughage Products, Sodium Sesquicarbonate, Potassium Chloride, Sodium Chloride and Processed Grain By-Products, Calcium Carbonate and Vitamin A Supplement, Carbohydrate (Glucose or corn starch), Gelatin, Ethoxyquine, Cane Molasses, Liquid Feed Syrup, Vegetable Fat (Feed Grade), Phosphoric acid, Soy Lecithin, Sulfuric Acid, Propionic Acid (A Preservative), Sorbic Acid (A Preservative), Benzoic Acid (A Preservative), Monocalcium Phosphate, Dicalcium Phosphate, Vitamin E Supplement, Selenium Yeast, Calcium Sulfate, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper, Amino Acid Complex, Cobalt Glucoheptonate complex, Magnesium Oxide, Manganese Sulfate, Zinc Sulfate, Potassium Chloride, Vitamin A Supplement, Vitamin D3 Supplement, Copper Sulfate, Ethylenediamine Dihydrate, Cobalt Carbonate, Diflubenzuron (CAS#35367-38-5), Montmorillonite Clay, Diatomaceous Earth, Hydrated Sodium Calcium Aluminosilicate, Yeast Brewers Dehydrated, Yeast Extract, Calcium Propionate (a preservative), Citric Acid (a preservative), BHT (a preservative), Silicon Dioxide, Vegetable Oil, Ammonium Hydroxide, Water, Natural and Artificial Flavoring.

FEEDING DIRECTIONS

Feed continuously at the rate of 1.25 pound of feed/100 pound of body weight; do not exceed 10.0 pounds per day. Feeding at 1.25 pound of feed/100 pound of body weight will supply Monensin at the level to control coccidiosis. Feed 18% Phoenix Feeds Calf Starter - R daily and supply fresh water at all times. Top quality hay may be introduced once dry feed intake reaches 5 pounds per day. When feed as directed this feed will provide sufficient Diflubenzuron Larvicide to prevent the emergence of adult house, stable, horn and face flies from the manure of treated cattle. Existing adult flies will not be affected. For optimal fly growth regulation, Diflubenzuron treated feed should first be fed 30 days prior to the last frost, continuously through the fly season and end 30 days after the first frost.

Caution: Do not allow horses or other equine access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin-medicated cattle and goat feeds are safe for use in cattle goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the feed refusals and the amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning: A withdrawal period has not been established for Monensin in pre-ruminating calves. Do not use in calves to be processed for veal.

03.07.24

Manufactured For
Phoenix Feeds, New Haven, VT 05472

Net Weight 50 pounds